

Charts, signatures, and stabilizations of Lefschetz fibrations

HISAAKI ENDO
ISAO HASEGAWA
SEIICHI KAMADA

We employ a certain labeled finite graph, called a chart, in a closed oriented surface for describing the monodromy of a(n achiral) Lefschetz fibration over the surface. Applying charts and their moves with respect to Wajnryb's presentation of mapping class groups, we first generalize a signature formula for Lefschetz fibrations over the 2-sphere obtained by Endo and Nagami to that for Lefschetz fibrations over arbitrary closed oriented surface. We then show two theorems on stabilization of Lefschetz fibrations under fiber summing with copies of a typical Lefschetz fibration as generalizations of a theorem of Auroux.

57M15; 57N13

1 Introduction

Matsumoto [28] proved that every Lefschetz fibration of genus one over a closed oriented surface is isomorphic to a fiber sum of copies of a holomorphic elliptic fibration on $\mathbb{CP}^2 \# 9\overline{\mathbb{CP}}^2$ and a trivial torus bundle over the surface if it has at least one critical point. This result played a crucial role in completing the classification of diffeomorphism types of elliptic surfaces (see Gompf and Stipsicz [11, Section 8.3]). Although such a classification has not been established for Lefschetz fibrations of higher genus, Auroux [1] proved a stabilization theorem for Lefschetz fibrations of genus two, which states that every Lefschetz fibration of genus two over the 2-sphere becomes isomorphic to a fiber sum of copies of three typical fibrations after fiber summing with a holomorphic fibration on $\mathbb{CP}^2 \# 13\overline{\mathbb{CP}}^2$. Auroux [2] gave a generalization of this theorem for Lefschetz fibrations of higher genus, which states that two Lefschetz fibrations of the same genus over the 2-sphere which have the same signature, the same numbers of singular fibers of each type, and admit sections of the same self-intersection number become isomorphic after fiber summing the same number of copies of a 'universal' Lefschetz fibration.

Kamada [15, 16] introduced charts, which are labeled finite graphs in a disk, to describe monodromies of surface braids (see also a textbook [17] of Kamada). Kamada, Matsumoto, Matumoto, and Waki [20] considered a variant of chart for Lefschetz fibrations of genus one to reprove the above result of Matsumoto. Furthermore Kamada [19], and Endo and Kamada [5, 6] made use of generalized charts to reprove the above theorem of Auroux for Lefschetz fibrations of genus two, and to investigate a stabilization theorem and an invariant for hyperelliptic Lefschetz fibrations of arbitrary genus. See also Baykur and Kamada [4], and Hayano [14] for applications of charts to broken Lefschetz fibrations.

In this paper we introduce a chart description for Lefschetz fibrations of genus greater than two over closed oriented surfaces of arbitrary genus to show a signature formula and two theorems on stabilization for such fibrations. In Section 2 we introduce charts and chart moves with respect to Wajnryb’s presentation of mapping class groups to examine monodromies of Lefschetz fibrations. After a short survey of Meyer’s signature cocycle, we generalize a signature formula [7] of Endo and Nagami for Lefschetz fibrations over the 2–sphere to that for Lefschetz fibrations over a closed oriented surface of arbitrary genus in Section 3. Section 4 is devoted to proofs of two theorems on stabilization of Lefschetz fibrations under fiber summing with copies of a ‘universal’ Lefschetz fibration. In particular the first of our stabilization theorem is a generalization of the theorem of Auroux [2]. We make several comments on variations of chart description and propose some possible directions for future research in Section 5.

2 Chart description for Lefschetz fibrations

In this section we review a definition and properties of Lefschetz fibrations and introduce a chart description for Lefschetz fibrations of genus greater than two.

2.1 Lefschetz fibrations and their monodromies

In this subsection we review a precise definition and basic properties of Lefschetz fibrations. More details can be found in Matsumoto [29] and Gompf and Stipsicz [11].

Let Σ_g be a closed oriented surface of genus g .

Definition 2.1 Let M and B be connected closed oriented smooth 4-manifold and 2-manifold, respectively. A smooth map $f: M \rightarrow B$ is called a *Lefschetz fibration* of genus g if it satisfies the following conditions:

- (i) the set $\Delta \subset B$ of critical values of f is finite and f is a smooth fiber bundle over $B - \Delta$ with fiber Σ_g ;
- (ii) for each $b \in \Delta$, there exists a unique critical point p in the *singular fiber* $F_b := f^{-1}(b)$ such that f is locally written as $f(z_1, z_2) = z_1 z_2$ or $\bar{z}_1 z_2$ with respect to some local complex coordinates around p and b which are compatible with orientations of M and B ;
- (iii) no fiber contains a (± 1) -sphere.

We call M the *total space*, B the *base space*, and f the *projection*. We call p a critical point of *positive type* (resp. of *negative type*) and F_b a singular fiber of *positive type* (resp. of *negative type*) if f is locally written as $f(z_1, z_2) = z_1 z_2$ (resp. $f(z_1, z_2) = \bar{z}_1 z_2$) in (ii). For a regular value $b \in B$ of f , $f^{-1}(b)$ is often called a *general fiber*.

Remark 1 A Lefschetz fibration in this paper is called an *achiral* Lefschetz fibration in many other papers.

Let $f: M \rightarrow B$ and $f': M' \rightarrow B$ be Lefschetz fibrations of genus g over the same base space B . We say that f is *isomorphic* to f' if there exist orientation preserving diffeomorphisms $H: M \rightarrow M'$ and $h: B \rightarrow B$ which satisfy $f' \circ H = h \circ f$. If we can choose such an h isotopic to the identity relative to a given base point $b_0 \in B$, we say that f is *strictly isomorphic* to f' .

Let $\mathcal{M}_g$ be the mapping class group of Σ_g , namely the group of all isotopy classes of orientation preserving diffeomorphisms of Σ_g . We assume that $\mathcal{M}_g$ acts on the *right*: the symbol $\varphi\psi$ means that we apply φ first and then ψ for $\varphi, \psi \in \mathcal{M}_g$. We denote the mapping class group of Σ_g acting on the left by $\mathcal{M}_g^*$. Hence the identity map $\mathcal{M}_g \rightarrow \mathcal{M}_g^*$ is an anti-isomorphism.

Let $f: M \rightarrow B$ be a Lefschetz fibration of genus g as in Definition 2.1. Take a base point $b_0 \in B$ and an orientation preserving diffeomorphism $\Phi: \Sigma_g \rightarrow F_0 := f^{-1}(b_0)$. Since f restricted over $B - \Delta$ is a smooth fiber bundle with fiber Σ_g , we can define a homomorphism

$$\rho: \pi_1(B - \Delta, b_0) \rightarrow \mathcal{M}_g$$

called the *monodromy representation* of f with respect to Φ . Let γ be the loop consisting of the boundary circle of a small disk neighborhood of $b \in \Delta$ oriented

counterclockwise and a simple path connecting a point on the circle to b_0 in $B - \Delta$. It is known that $\rho(\gamma)$ is a Dehn twist along some essential simple closed curve c on Σ_g , which is called the *vanishing cycle* of the critical point p on $f^{-1}(b)$. If p is of positive type (resp. of negative type), then the Dehn twist is right-handed (resp. left-handed).

A singular fiber is said to be of *type I* if the vanishing cycle is non-separating and of *type II_h* for $h = 1, \dots, [g/2]$ if the vanishing cycle is separating and it bounds a genus- h subsurface of Σ_g . A singular fiber is said to be of *type I⁺* (resp. *type I⁻* and *type II_h⁺*, *type II_h⁻*) if it is of type I and of positive type (resp. of type I and of negative type, of type II_h and of positive type, of type II_h and of negative type). We denote by $n_0^+(f)$, $n_0^-(f)$, $n_h^+(f)$, and $n_h^-(f)$, the numbers of singular fibers of f of type I⁺, I⁻, II_h⁺, and II_h⁻, respectively. A Lefschetz fibration is called *irreducible* if every singular fiber is of type I. A Lefschetz fibration is called *chiral* if every singular fiber is of positive type.

Suppose that the cardinality of Δ is equal to n . A system $\mathcal{A} = (A_1, \dots, A_n)$ of arcs on B is called a *Hurwitz arc system* for Δ with base point b_0 if each A_i is an embedded arc connecting b_0 with a point of Δ in B such that $A_i \cap A_j = \{b_0\}$ for $i \neq j$, and they appear in this order around b_0 (see Kamada [17]). When B is a 2-sphere, the system $\mathcal{A}$ determines a system of generators of $\pi_1(B - \Delta, b_0)$, say $(a_1, \dots, a_n)$. We call $(\rho(a_1), \dots, \rho(a_n))$ a *Hurwitz system* of f .

2.2 Chart description and Wajnryb's presentation


In this subsection we introduce a chart description for Lefschetz fibrations of genus greater than two by employing Wajnryb's finite presentation [36] of mapping class groups. We use the terminology of chart description in Kamada [18].

We first review a finite presentation of the mapping class group of a closed oriented surface due to Wajnryb. For $i = 0, 1, \dots, 2g$, let ζ_i be a right-handed Dehn twist along the simple closed curve c_i on Σ_g depicted in Figure 1.

Theorem 2.2 (Wajnryb [36, 37]) *Suppose that g is greater than two. The mapping class group $\mathcal{M}_g$ is generated by elements $\zeta_0, \zeta_1, \zeta_2, \dots, \zeta_{2g}$ and has defining relations:*

$$(A) \quad \begin{aligned} \zeta_i \zeta_j &= \zeta_j \zeta_i \quad (1 \leq i < j - 1 \leq 2g - 1), \quad \zeta_0 \zeta_j = \zeta_j \zeta_0 \quad (j = 1, 2, 3, 5, \dots, 2g), \\ \zeta_i \zeta_{i+1} \zeta_i &= \zeta_{i+1} \zeta_i \zeta_{i+1} \quad (i = 1, \dots, 2g - 1), \quad \zeta_0 \zeta_4 \zeta_0 = \zeta_4 \zeta_0 \zeta_4; \end{aligned}$$

$$(B) \quad (\zeta_3 \zeta_2 \zeta_1)^4 = \zeta_0 \zeta_4^{-1} \zeta_3^{-1} \zeta_2^{-1} \zeta_1^{-2} \zeta_2^{-1} \zeta_3^{-1} \zeta_4^{-1} \zeta_0 \zeta_4 \zeta_3 \zeta_2 \zeta_1^2 \zeta_2 \zeta_3 \zeta_4;$$

Figure 1: Simple closed curves on Σ_g

$$(C) \quad \delta_3 \zeta_1 \zeta_3 \zeta_5 = \zeta_0 \tau_2 \zeta_0 \tau_2^{-1} \tau_1 \tau_2 \zeta_0 \tau_2^{-1} \tau_1^{-1}, \quad \text{where}$$

$$\begin{aligned} \tau_1 &:= \zeta_2 \zeta_3 \zeta_1 \zeta_2, & \tau_2 &:= \zeta_4 \zeta_5 \zeta_3 \zeta_4, & \mu &:= \zeta_5 \zeta_6 \tau_2 \zeta_0 \tau_2^{-1} \zeta_6^{-1} \zeta_5^{-1}, \\ \nu &:= \zeta_1 \zeta_2 \zeta_3 \zeta_4 \zeta_0 \zeta_1^{-1} \zeta_2^{-1} \zeta_3^{-1} \zeta_4^{-1}, & \delta_3 &:= \zeta_6^{-1} \zeta_5^{-1} \zeta_4^{-1} \zeta_3^{-1} \zeta_2^{-1} \mu^{-1} \nu \mu \zeta_2 \zeta_3 \zeta_4 \zeta_5 \zeta_6; \end{aligned}$$

$$(D) \quad \zeta_{2g} \cdots \zeta_3 \zeta_2 \zeta_1^2 \zeta_2 \zeta_3 \cdots \zeta_{2g} \delta_g = \delta_g \zeta_{2g} \cdots \zeta_3 \zeta_2 \zeta_1^2 \zeta_2 \zeta_3 \cdots \zeta_{2g}, \quad \text{where}$$

$$\begin{aligned} \tau_1 &:= \zeta_2 \zeta_3 \zeta_1 \zeta_2, & \tau_i &:= \zeta_{2i} \zeta_{2i-1} \zeta_{2i+1} \zeta_{2i}, \\ \nu_1 &:= \zeta_4^{-1} \zeta_3^{-1} \zeta_2^{-1} \zeta_1^{-2} \zeta_2^{-1} \zeta_3^{-1} \zeta_4^{-1} \zeta_0 \zeta_4 \zeta_3 \zeta_2 \zeta_1^2 \zeta_2 \zeta_3 \zeta_4, & \nu_i &:= \tau_{i-1} \tau_i \nu_{i-1} \tau_i^{-1} \tau_{i-1}^{-1}, \\ \mu_1 &:= \zeta_2 \zeta_3 \zeta_4 \nu_1 \zeta_1^{-1} \zeta_2^{-1} \zeta_3^{-1} \zeta_4^{-1}, & \mu_i &:= \zeta_{2i} \zeta_{2i+1} \zeta_{2i+2} \nu_i \zeta_{2i-1}^{-1} \zeta_{2i}^{-1} \zeta_{2i+1}^{-1} \zeta_{2i+2}^{-1}, \\ \delta_g &:= \mu_{g-1}^{-1} \cdots \mu_2^{-1} \mu_1^{-1} \zeta_1 \mu_1 \mu_2 \cdots \mu_{g-1} \end{aligned}$$

for $i = 2, \dots, g-1$.

We make use of the presentation above to introduce a notion of chart which gives a graphic description of monodromy representations of Lefschetz fibrations. We set

$$\begin{aligned} \mathcal{X} &:= \{\zeta_0, \zeta_1, \dots, \zeta_{2g}\}, \\ \mathcal{R} &:= \{r_1(i, j) \mid 1 \leq i < j-1 \leq 2g-1\} \cup \{r_1(0, j) \mid j = 1, 2, 3, 5, \dots, 2g\} \\ &\quad \cup \{r_2(i) \mid i = 0, 1, \dots, 2g-1\} \cup \{r_3, r_4, r_5\}, \\ \mathcal{S} &:= \{\ell_1(i)^{\pm 1} \mid i = 0, 1, \dots, 2g\} \cup \{\ell_2(h)^{\pm 1} \mid h = 1, \dots, [g/2]\}, \end{aligned}$$

for $g \geq 3$, where

$$\begin{aligned} r_1(i, j) &:= \zeta_i \zeta_i \zeta_i^{-1} \zeta_j^{-1}, & r_2(0) &:= \zeta_0 \zeta_4 \zeta_0 \zeta_4^{-1} \zeta_0^{-1} \zeta_4, \\ r_2(i) &:= \zeta_i \zeta_{i+1} \zeta_i \zeta_{i+1}^{-1} \zeta_i^{-1} \zeta_{i+1} \quad (i = 1, \dots, 2g-1), \end{aligned}$$

$$\begin{aligned}
r_3 &:= (\zeta_3 \zeta_2 \zeta_1)^4 \zeta_4^{-1} \zeta_3^{-1} \zeta_2^{-1} \zeta_1^{-2} \zeta_2^{-1} \zeta_3^{-1} \zeta_4^{-1} \zeta_0^{-1} \zeta_4 \zeta_3 \zeta_2 \zeta_1^2 \zeta_2 \zeta_3 \zeta_4 \zeta_0^{-1}, \\
r_4 &:= \delta_3 \zeta_1 \zeta_3 \zeta_5 \tau_1 \tau_2 \zeta_0^{-1} \tau_2^{-1} \tau_1^{-1} \tau_2 \zeta_0^{-1} \tau_2^{-1} \zeta_0^{-1}, \\
r_5 &:= \zeta_{2g} \cdots \zeta_3 \zeta_2 \zeta_1^2 \zeta_2 \zeta_3 \cdots \zeta_{2g} \delta_g \zeta_{2g}^{-1} \cdots \zeta_3^{-1} \zeta_2^{-1} \zeta_1^{-2} \zeta_2^{-1} \zeta_3^{-1} \cdots \zeta_{2g}^{-1} \delta_g^{-1}, \\
\ell_1(i) &:= \zeta_i \quad (i = 0, 1, \dots, 2g), \quad \ell_2(h) := (\zeta_1 \zeta_2 \cdots \zeta_{2h})^{4h+2} \quad (h = 1, \dots, [g/2]),
\end{aligned}$$

and $\delta_3, \tau_1, \tau_2, \delta_g$ are defined as in Theorem 2.2.

Let B be a connected closed oriented surface and Γ a finite graph in B such that each edge of Γ is oriented and labeled with an element of $\mathcal{X}$. For a vertex v of Γ , a small simple closed curve surrounding v in the positive direction of B is called a *meridian loop* of v and denoted by m_v . The vertex v is said to be *marked* if one of the regions around v is specified by an asterisk. If v is marked, the intersection word $w_\Gamma(m_v)$ of m_v with respect to Γ is well-defined. If not, it is determined up to cyclic permutation. See Kamada [18] for details.

Definition 2.3 A *chart* in B is a finite graph Γ in B (possibly being empty or having *hoops* that are closed edges without vertices) whose edges are labeled with an element of $\mathcal{X}$, and oriented so that the following conditions are satisfied (see Figure 2, Figure 3, and Figure 4):

- (1) the vertices of Γ are classified into two families: *white vertices* and *black vertices*;
- (2) if v is a white vertex (resp. a black vertex), the word $w_\Gamma(m_v)$ is a cyclic permutation of an element of $\mathcal{R} \cup \mathcal{R}^{-1}$ (resp. of $\mathcal{S}$).

A white vertex v is said to be of *type* r (resp. of *type* r^{-1}) if $w_\Gamma(m_v)^{-1}$ is a cyclic permutation of $r \in \mathcal{R}$ (resp. of $r^{-1} \in \mathcal{R}^{-1}$). A black vertex v is said to be of *type* s if $w_\Gamma(m_v)$ is a cyclic permutation of $s \in \mathcal{S}$. A chart Γ is said to be *marked* if each white vertex (resp. black vertex) v is marked and $w_\Gamma(m_v)$ is exactly an element of $\mathcal{R} \cup \mathcal{R}^{-1}$ (resp. of $\mathcal{S}$). If a base point b_0 of B is specified, we always assume that a chart Γ is disjoint from b_0 . A chart consisting of two black vertices and one edge connecting them is called a *free edge*. We denote the label ζ_i by i for short.






Figure 2: Vertices of type $\ell_1(i)^{\pm 1}$, $r_1(i, j)$, $r_2(i)$ ($i \neq 0$)

We next introduce several moves for charts. Let Γ and Γ' be two charts on B and b_0 a base point of B .

Figure 3: Vertices of type r_3 and r_4 Figure 4: Vertices of type $\ell_2(h)$ and r_5

Let D be a disk embedded in $B - \{b_0\}$. Suppose that the boundary ∂D of D intersects Γ and Γ' transversely.

Definition 2.4 We say that Γ' is obtained from Γ by a *chart move of type W* if $\Gamma \cap (B - \text{Int } D) = \Gamma' \cap (B - \text{Int } D)$ and that both $\Gamma \cap D$ and $\Gamma' \cap D$ have no black vertices. We call chart moves of type W shown in Figure 5 (a), (b), and (c), a *channel change*, a *birth/death of a hoop*, and a *birth/death of a pair of white vertices*, respectively.




Figure 5: Chart moves of type W

Let s and s' be words in $\mathcal{S}$. Suppose that there exists a word w in $\mathcal{X} \cup \mathcal{X}^{-1}$ such that two words s' and $ws w^{-1}$ determine the same element of $\mathcal{M}_g$.

Definition 2.5 If a chart Γ contains a black vertex of type s , then we can change a part of Γ near the vertex by using a local replacement depicted in Figure 6 to obtain

another chart Γ' . We say that Γ' is obtained from Γ by a *chart move of transition*. Note that the blank labeled with T can be filled only with edges and white vertices.




Figure 6: Chart move of transition

Definition 2.6 We say that Γ' is obtained from Γ by a *chart move of conjugacy type* if Γ' is obtained from Γ by a local replacement depicted in Figure 7.




Figure 7: Chart moves of conjugacy type

Let Γ be a chart in B with base point b_0 and Δ_Γ the set of black vertices of Γ . For a loop η in $B - \Delta_\Gamma$ based at b_0 , the element of $\mathcal{M}_g$ determined by the intersection word $w_\Gamma(\eta)$ of η with respect to Γ does not depend on a choice of representative of the homotopy class of η . Thus we obtain a homomorphism $\rho_\Gamma: \pi_1(B - \Delta_\Gamma, b_0) \rightarrow \mathcal{M}_g$, which is called the *homomorphism determined by Γ* .

We now state a classification of Lefschetz fibrations in terms of charts and chart moves. Let B be a connected closed oriented surface.

Proposition 2.7 Suppose that g is greater than two. (1) Let f be a Lefschetz fibration of genus g over B and ρ a monodromy representation of f . Then there exists a chart Γ in B such that the homomorphism ρ_Γ determined by Γ is equal to ρ . (2) For every chart Γ in B , there exists a Lefschetz fibration f of genus g over B such that a monodromy representation of f is equal to the homomorphism ρ_Γ determined by Γ .

We call such Γ as in Proposition 2.7 (1) a chart *corresponding to f* , and such f as in Proposition 2.7 (2) a Lefschetz fibration *described by Γ* .

Theorem 2.8 Suppose that g is greater than two. Let f and f' be Lefschetz fibrations of genus g over B , and Γ and Γ' charts corresponding to f and f' , respectively. Then f is strictly isomorphic to f' if and only if Γ is transformed to Γ' by a finite sequence of chart moves of type W , chart moves of transitions, chart moves of conjugacy type, and ambient isotopies of B relative to b_0 .

Proofs of Proposition 2.7 and Theorem 2.8 are straightforward from a classification theorem of Lefschetz fibrations due to Kas [21] and Matsumoto [29] together with fundamental theorems on charts and chart moves by Kamada [18, Sections 4–8].

We end this subsection with a definition and chart description of fiber sums of Lefschetz fibrations. Let $f: M \rightarrow B$ and $f': M' \rightarrow B'$ be Lefschetz fibrations of genus g . Take regular values $b_0 \in B$ and $b'_0 \in B'$ of f and f' , and small disks $D_0 \subset B - \Delta$ and $D'_0 \subset B' - \Delta'$ near b_0 and b'_0 , respectively. Consider general fibers $F_0 := f^{-1}(b_0)$ and $F'_0 := f'^{-1}(b'_0)$ and orientation preserving diffeomorphisms $\Phi: \Sigma_g \rightarrow F_0$ and $\Phi': \Sigma_g \rightarrow F'_0$, respectively.

Definition 2.9 Let $\Psi: \Sigma_g \rightarrow \Sigma_g$ be an orientation preserving diffeomorphism and $r: \partial D_0 \rightarrow \partial D'_0$ an orientation reversing diffeomorphism. The new manifold $M \#_F M'$ obtained by glueing $M - f^{-1}(\text{Int } D_0)$ and $M' - f'^{-1}(\text{Int } D'_0)$ by $(\Phi' \circ \Psi \circ \Phi^{-1}) \times r$ admits a Lefschetz fibration $f \#_\Psi f': M \#_F M' \rightarrow B \# B'$ of genus g . We call $f \#_\Psi f'$ the *fiber sum* of f and f' with respect to Ψ . Although the diffeomorphism type of $M \#_F M'$ and the isomorphism type of $f \#_\Psi f'$ depend on a choice of the diffeomorphism Ψ in general, we often abbreviate $f \#_\Psi f'$ as $f \# f'$.

Let Γ and Γ' be charts corresponding to f and f' , and D_0 and D'_0 small disks near b_0 and b'_0 disjoint from Γ and Γ' , respectively. Connecting $B - \text{Int } D_0$ with $B' - \text{Int } D'_0$ by a tube, we have a connected sum $B \# B'$ of B and B' . Let w be a word in $\mathcal{X} \cup \mathcal{X}^{-1}$ which represents the mapping class of Ψ in $\mathcal{M}_g$. Let $\Gamma \#_w \Gamma'$ be the union of Γ , Γ' , and hoops on the tube representing w (see Figure 8). Then the fiber sum $f \#_\Psi f'$ is described by this new chart $\Gamma \#_w \Gamma'$ in $B \# B'$ with base point b_0 . If the word w is trivial, then the chart $\Gamma \#_w \Gamma'$ is denoted also by $\Gamma \oplus \Gamma'$, which is called a *product* of Γ and Γ' .




Figure 8: Chart $\Gamma \#_w \Gamma'$ in $B \# B'$

3 Signature of Lefschetz fibrations

In this section we review the signature cocycle discovered by Meyer and prove a signature theorem for Lefschetz fibrations.

3.1 Meyer's signature cocycle

In this subsection we give a brief survey on Meyer's signature cocycle. We begin with the definition of the signature cocycle. Let g be a positive integer.

Definition 3.1 (Meyer [30]) For $A, B \in \mathrm{Sp}(2g, \mathbb{Z})$, we consider the vector space

$$V_{A,B} := \{(x, y) \in \mathbb{R}^{2g} \times \mathbb{R}^{2g} \mid (A^{-1} - I_{2g})x + (B - I_{2g})y = 0\}$$

and the bilinear form $\langle \cdot, \cdot \rangle_{A,B}: V_{A,B} \times V_{A,B} \rightarrow \mathbb{R}$ defined by

$$\langle (x_1, y_1), (x_2, y_2) \rangle_{A,B} := (x_1 + y_1) \cdot J(I_{2g} - B)y_2,$$

where $\cdot$ is the standard inner product of $\mathbb{R}^{2g}$ and $J = \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix}$. Since $\langle \cdot, \cdot \rangle_{A,B}$ is symmetric, we can define an integer $\tau_g(A, B)$ to be the signature of $(V_{A,B}, \langle \cdot, \cdot \rangle_{A,B})$. The map $\tau_g: \mathrm{Sp}(2g, \mathbb{Z}) \times \mathrm{Sp}(2g, \mathbb{Z}) \rightarrow \mathbb{Z}$ is called the *signature cocycle*.

Let P be a compact connected oriented surface of genus 0 with three boundary components and $\pi: E \rightarrow P$ a fiber bundle over P with fiber Σ_g and structure group $\mathrm{Diff}_+\Sigma_g$. The fundamental group $\pi_1(P, *)$ of P with base point $*$ is a free group generated by two loops a and b depicted in Figure 9. If we take an orientation preserving diffeomorphism $\Sigma_g \rightarrow \pi^{-1}(*)$, we obtain the monodromy representation $\pi_1(P, *) \rightarrow \mathcal{M}_g$ which sends a to α and b to β . Since $\mathcal{M}_g^*$ acts on $H := H_1(\Sigma_g; \mathbb{Z})$ preserving the intersection form, we have a representation $\mathcal{M}_g^* \rightarrow \mathrm{Sp}(2g, \mathbb{Z})$ by fixing a symplectic basis on H . Let A and B denote matrices corresponding to α and β , respectively.




Figure 9: Pair of pants P

Meyer closely studied the signature of the total space E to obtain the following theorem.

Theorem 3.2 (Meyer [30]) *The signature $\sigma(E)$ of E is equal to $-\tau_g(A, B)$.*

Theorem 3.2 and Novikov's additivity implies that τ_g is a 2-cocycle of $\mathrm{Sp}(2g, \mathbb{Z})$.

We recall a Maslov index for a triple of Lagrangian subspaces and Wall's non-additivity theorem, which are used in the proof of Theorem 3.2.

Let V be a real vector space of dimension $2n$, $\omega \in \Lambda^2 V^*$ a symplectic form on V , and $\Lambda(V, \omega)$ the Lagrangian Grassmannian of (V, ω) , which is the set of Lagrangian subspaces of (V, ω) . For $L_1, L_2, L_3 \in \Lambda(V, \omega)$, the bilinear form

$$\begin{aligned} \Psi: (L_3 + L_1) \cap L_2 \times (L_3 + L_1) \cap L_2 &\rightarrow \mathbb{R}: (v, w) \mapsto \omega(v, w_3) \\ (v, w \in (L_3 + L_1) \cap L_2, w &= w_1 + w_3 (w_1 \in L_1, w_3 \in L_3)) \end{aligned}$$

is symmetric. We define an integer $i(L_1, L_2, L_3)$ to be the signature of $((L_3 + L_1) \cap L_2, \Psi)$, which is called the *ternary Maslov index* of the triple (L_1, L_2, L_3) .

Let M_1, M_2 be compact oriented smooth 4-manifolds, X_1, X_2, X_3 compact oriented smooth 3-manifolds, and Σ a closed oriented smooth 2-manifold. We assume that $M = M_1 \cup M_2$, $\partial M_1 = X_1 \cup X_2$, $\partial M_2 = X_2 \cup X_3$, $\partial X_1 = \partial X_2 = \partial X_3 = \Sigma$, and the orientations of these manifolds satisfy

$$\begin{aligned} [M] &= [M_1] + [M_2], \quad \partial_*[M_1] = [X_2] - [X_1], \quad \partial_*[M_2] = [X_3] - [X_2], \\ \partial_*[X_1] &= \partial_*[X_2] = \partial_*[X_3] = [\Sigma]. \end{aligned}$$

Let $\omega: V \times V \rightarrow \mathbb{R}$ be the intersection form on $V := H_1(\Sigma; \mathbb{R})$ and L_i the kernel of the homomorphism $V \rightarrow H_1(X_i; \mathbb{R})$ induced by the inclusion $\Sigma \rightarrow X_i$ for $i = 1, 2, 3$. Since $L_i \in \Lambda(V, \omega)$ for $i = 1, 2, 3$, we can define the ternary Maslov index $i(L_1, L_2, L_3)$ of the triple (L_1, L_2, L_3) .

Theorem 3.3 (Wall [38]) $\sigma(M) = \sigma(M_1) + \sigma(M_2) - i(L_1, L_2, L_3)$.

Gambaudo and Ghys [9] (and independently the first author) made use of Theorem 3.3 to give the following proof of Theorem 3.2.

Proof of Theorem 3.2 Consider P to be a boundary sum of two annuli P_1 and P_2 (see Figure 9). We set $M := E$, $M_i := \pi^{-1}(P_i)$ ($i = 1, 2$), $X_2 := M_1 \cap M_2$, $X_1 := \partial M_1 - \mathrm{Int} X_2$, $X_3 := \partial M_2 - \mathrm{Int} X_2$, and $\Sigma := \partial X_2$. Applying Theorem 3.3 to these manifolds, we have

$$\sigma(E) = \sigma(M_1) + \sigma(M_2) - i(L_1, L_2, L_3) = -i(L_1, L_2, L_3)$$

because each of M_1 and M_2 is a product of a mapping torus with an interval, which has signature zero. Since the bordered component of X_i is diffeomorphic to $I \times \Sigma_g$ for

$i = 1, 2, 3$, we put $V := H \oplus H$, $\omega := \mu \oplus (-\mu)$, and obtain

$$\begin{aligned} L_1 &= \{(-\xi, \alpha_*^{-1}(\xi)) \in V \mid \xi \in H\}, \quad L_2 = \{(-\xi, \xi) \in V \mid \xi \in H\}, \\ L_3 &= \{(-\xi, \beta_*(\xi)) \in V \mid \xi \in H\}, \end{aligned}$$

where H is the first homology $H_1(\Sigma_g; \mathbb{R})$ of Σ_g and $\mu: H \times H \rightarrow \mathbb{R}$ is the intersection form of Σ_g . It is easily seen that the subspace $(L_1 + L_3) \cap L_2$ is written as

$$(L_1 + L_3) \cap L_2 = \{(-\xi - \eta, \alpha_*^{-1}(\xi) + \beta_*(\eta)) \in V \mid \xi + \eta = \alpha_*^{-1}(\xi) + \beta_*(\eta) \ (\xi, \eta \in H)\}$$

and the symmetric bilinear form Ψ on $(L_1 + L_3) \cap L_2$ is written as

$$\Psi((-\xi - \eta, \alpha_*^{-1}(\xi) + \beta_*(\eta)), (-\xi' - \eta', \alpha_*^{-1}(\xi') + \beta_*(\eta'))) = \mu(\xi + \eta, (\text{id} - \beta_*)(\eta')).$$

We consider the vector space

$$U_{\alpha, \beta} := \{(\xi, \eta) \in V \mid (\alpha_*^{-1} - \text{id})(\xi) + (\beta_* - \text{id})(\eta) = 0\}$$

and the symmetric bilinear form $\langle \cdot, \cdot \rangle_{\alpha, \beta}$ on $U_{\alpha, \beta}$ defined by

$$\langle (\xi, \eta), (\xi', \eta') \rangle_{\alpha, \beta} := \mu(\xi + \eta, (\text{id} - \beta_*)(\eta')) \quad ((\xi, \eta), (\xi', \eta') \in U_{\alpha, \beta}).$$

Since the linear map $U_{\alpha, \beta} \rightarrow (L_1 + L_3) \cap L_2: (\xi, \eta) \mapsto (-\xi - \eta, \xi + \eta)$ is compatible with the bilinear forms, the signature of $((L_1 + L_3) \cap L_2, \Psi)$ is equal to that of $(U_{\alpha, \beta}, \langle \cdot, \cdot \rangle_{\alpha, \beta})$, which is isomorphic to $(V_{A, B}, \langle \cdot, \cdot \rangle_{A, B})$ under a choice of a symplectic basis of H . Therefore we conclude that $i(L_1, L_2, L_3) = \tau_g(A, B)$. $\square$

Remark 2 It is known that τ_g is a normalized, symmetric 2-cocycle of $\text{Sp}(2g, \mathbb{Z})$ and invariant under conjugation. The cohomology class $[\tau_g] \in H^2(\text{Sp}(2g, \mathbb{Z}); \mathbb{Z})$ corresponds to $-4c_1$ under homomorphisms:

$$H^2(\text{Sp}(2g, \mathbb{Z}); \mathbb{Z}) \leftarrow H^2(\text{BSp}(2g, \mathbb{R}); \mathbb{Z}) \cong H^2(\text{BU}(g); \mathbb{Z}) \cong \mathbb{Z}.$$

For more details see Meyer [30], Turaev [35], Barge and Ghys [3], and Kuno [24].

3.2 A signature formula

In this subsection we describe the signature of a Lefschetz fibration of genus greater than two in terms of charts. Let g be an integer greater than two.

Let B be a connected closed oriented surface and Γ a chart in B . We denote the number of white vertices of type $r_1(i, j)$ (resp. $r_2(i), r_3, r_4, r_5$) minus the number of white vertices of type $r_1(i, j)^{-1}$ (resp. $r_2(i)^{-1}, r_3^{-1}, r_4^{-1}, r_5^{-1}$) included in Γ by $n_1(i, j)(\Gamma)$ (resp. $n_2(i)(\Gamma), n_3(\Gamma), n_4(\Gamma), n_5(\Gamma)$). Similarly, we denote the number of black vertices of type $\ell_1(i)^{\pm 1}$ (resp. $\ell_2(h)^{\pm 1}$) included in Γ by $m_1^{\pm}(i)(\Gamma)$ (resp. $m_2^{\pm}(h)(\Gamma)$), and set $m_1(i)(\Gamma) := m_1^+(i)(\Gamma) - m_1^-(i)(\Gamma)$ (resp. $m_2(h)(\Gamma) := m_2^+(h)(\Gamma) - m_2^-(h)(\Gamma)$) and $m_1^{\pm}(\Gamma) := \sum_{i=0}^{2g} m_1^{\pm}(i)(\Gamma)$.

Definition 3.4 The number

$$\sigma(\Gamma) := -6n_3(\Gamma) - n_4(\Gamma) + \sum_{h=1}^{\lfloor g/2 \rfloor} (4h(h+1) - 1)m_2(h)(\Gamma)$$

is called the *signature* of Γ .

Let $f: M \rightarrow B$ be a Lefschetz fibration of genus g and Γ a chart in B corresponding to f . The purpose of this subsection is to show the following theorem.

Theorem 3.5 *The signature $\sigma(M)$ of M is equal to $\sigma(\Gamma)$.*

Remark 3 It immediately follows from Theorem 3.5 that $\sigma(\Gamma)$ is invariant under chart moves of type W and chart moves of transition. Any combinatorial proof of this fact does not seem to be known.

Let $\tilde{\mathcal{X}}$ be the set of right-handed Dehn twists along simple closed curves in Σ_g and $\tilde{\mathcal{R}}$ the set of words in $\tilde{\mathcal{X}} \cup \tilde{\mathcal{X}}^{-1}$ representing an element of the kernel of the natural epimorphism from the free group generated by $\tilde{\mathcal{X}}$ to $\mathcal{M}_g$.

Definition 3.6 For a word $w = \alpha_1 \cdots \alpha_n \in \tilde{\mathcal{R}}$, we define an integer


$$I_g(w) := - \sum_{j=1}^{n-1} \tau_g(\overline{\alpha_{n-j}}, \overline{\alpha_{n-j+1}} \cdots \overline{\alpha_n}) - s(w),$$

where τ_g is the signature cocycle (Definition 3.1), $\overline{\alpha}$ is the image of $\alpha \in \tilde{\mathcal{X}} \cup \tilde{\mathcal{X}}^{-1}$ under the composition of the natural map $\tilde{\mathcal{X}} \cup \tilde{\mathcal{X}}^{-1} \rightarrow \mathcal{M}_g$ and a natural epimorphism $\mathcal{M}_g^* \rightarrow \text{Sp}(2g, \mathbb{Z})$, and $s(w)$ is the number of Dehn twists along separating simple closed curves included in w .

Suppose that B is a 2-sphere. If we choose a monodromy representation ρ and a Hurwitz arc system $\mathcal{A}$ for Δ with base point b_0 , we have a Hurwitz system $(\alpha_1, \dots, \alpha_n) \in (\mathcal{M}_g)^n$ of f . Since $\alpha_1, \dots, \alpha_n$ are Dehn twists and $\alpha_1 \cdots \alpha_n = 1$ in $\mathcal{M}_g$, we think $(\alpha_1, \dots, \alpha_n)$ as a word $w := \alpha_1 \cdots \alpha_n$ in $\tilde{\mathcal{R}}$. Theorem 3.2 and Novikov's additivity for signature imply the next theorem.

Theorem 3.7 (Endo and Nagami [7]) *The signature $\sigma(M)$ of M is equal to $I_g(w)$.*

We are now ready to prove Theorem 3.5.

Figure 10: A normal form of Γ

Proof of Theorem 3.5 Case 1: Suppose that B is a 2-sphere. We can assume that Γ is included in a disk D disjoint from a base point b_0 in B . Deforming Γ by an isotopy of D , we can make Γ in the form depicted in Figure 10, where the blank labeled with T is filled only with edges (and hoops). Let w_1 and w_2 be the intersection words of paths in D depicted in Figure 10 with respect to Γ .

It is easy to see that w_1 is written as

$$w_1 = u_1^{-1} \rho_1^{\varepsilon_1} u_1 \cdots u_r^{-1} \rho_r^{\varepsilon_r} u_r,$$

where $u_1, \dots, u_r$ are words in $\mathcal{X} \cup \mathcal{X}^{-1}$, $\rho_1, \dots, \rho_r \in \mathcal{R}$, and $\varepsilon_1, \dots, \varepsilon_r \in \{\pm 1\}$, and w_2 is written as

$$w_2 = v_1^{-1} \lambda_1 v_1 \cdots v_s^{-1} \lambda_s v_s,$$

where $v_1, \dots, v_s$ are words in $\mathcal{X} \cup \mathcal{X}^{-1}$ and $\lambda_1, \dots, \lambda_s \in \mathcal{S}$.

Let σ_h be a right-handed Dehn twist along the curve s_h depicted in Figure 1. We define two words w_3 and w_4 as follows. We set

$$w_3 := v_1^{-1} \tilde{\lambda}_1 v_1 \cdots v_s^{-1} \tilde{\lambda}_s v_s,$$

where $\tilde{\lambda}_j$ is equal to $\zeta_i^{\pm 1}$ (resp. $\sigma_h^{\pm 1}$) if λ_j is equal to $\ell_1(i)^{\pm 1}$ (resp. $\ell_2(h)^{\pm 1}$). For each $j = 1, \dots, s$, the word $v_j^{-1} \tilde{\lambda}_j v_j$ represents a Dehn twist $\xi_j \in \mathcal{X} \cup \mathcal{X}^{-1}$ because it is conjugate to the Dehn twist $\tilde{\lambda}_j$. We set $w_4 := \xi_1 \cdots \xi_s$.

Since all four words w_1, w_2, w_3, w_4 belong to $\tilde{\mathcal{R}}$, we can consider the values of the function $I_g: \tilde{\mathcal{R}} \rightarrow \mathbb{Z}$ for these words. Since $(\xi_1, \dots, \xi_s)$ is a Hurwitz system of f by

Proposition 2.7 (2), the signature $\sigma(M)$ of M is equal to $I_g(w_4)$ from Theorem 3.7. According to basic properties and explicit computations for I_g due to Endo and Nagami [7, Lemma 3.5, Proposition 3.6, Proposition 3.9], we have

$$\begin{aligned} I_g(w_4) &= I_g(w_3) = I_g(w_2) + \sum_{h=1}^{\lfloor g/2 \rfloor} m_2(h)(\Gamma) \cdot I_g((\zeta_1 \zeta_2 \cdots \zeta_{2h})^{-(4h+2)} \sigma_h) \\ &= I_g(w_2) + \sum_{h=1}^{\lfloor g/2 \rfloor} (4h(h+1) - 1) m_2(h)(\Gamma). \end{aligned}$$

On the other hand, we have also

$$\begin{aligned} I_g(w_1) &= I_g(u_1^{-1} \rho_1^{\varepsilon_1} u_1 \cdots u_r^{-1} \rho_r^{\varepsilon_r} u_r) = \varepsilon_1 I_g(\rho_1) + \cdots + \varepsilon_r I_g(\rho_r) \\ &= \sum_{i,j} n_1(i,j)(\Gamma) \cdot I_g(r_1(i,j)) + \sum_{i=0}^{2g-1} n_2(i)(\Gamma) \cdot I_g(r_2(i)) \\ &\quad + n_3(\Gamma) \cdot I_g(r_3) + n_4(\Gamma) \cdot I_g(r_4) + n_5(\Gamma) \cdot I_g(r_5) \\ &= -6n_3(\Gamma) - n_4(\Gamma) \end{aligned}$$

from [7, Lemma 3.5, Remark 3.7, Propositions 3.9 and 3.10]. Since w_1 is freely equal to w_2 as words in $\tilde{\mathcal{X}} \cup \tilde{\mathcal{X}}^{-1}$ and thus $I_g(w_1) = I_g(w_2)$, we obtain $\sigma(M) = \sigma(\Gamma)$.

Case 2: Suppose that B is not a 2-sphere. Then the genus g' of B is positive. Let b_0 be a base point of B and D a disk in $B - \Gamma$ centered at b_0 . We take a point $b_1 \in B - (\Gamma \cup D)$ and the union A of $2g'$ simple loops based at b_1 . Suppose that any two loops of A intersect only at b_1 and $B - A$ is an open disk (an open $4g'$ -gon). We also suppose that each loop of A intersects with edges of Γ transversely. For each $b \in \Gamma \cap A$, we choose a simple path γ_b from b to b_0 which intersects with edges of Γ transversely and does not intersect with $\Gamma \cap A$ other than b . Let w_b be the intersection word of γ_b with respect to Γ and $i_b \in \{0, 1, \dots, 2g\}$ the label of the edge including b . We choose a family $\{D_b\}_{b \in \Gamma \cap A}$ of disjoint disks included in D and put the chart Γ_b depicted in Figure 11 in D_b for each $b \in \Gamma \cap A$. Taking the union of Γ with Γ_b for all $b \in \Gamma \cap A$, we obtain a new chart Γ_1 in B . A Lefschetz fibration $f_1: M_1 \rightarrow B$ described by Γ_1 is nothing but a fiber sum of f with Lefschetz fibrations over S^2 described by a free edge.

For each $b \in \Gamma \cap A$, we apply channel changes as in Figure 12 to let a free edge pass through the edges intersecting with γ_b . We then apply a channel change as in Figure 13 to remove the intersection of Γ with A at b . Thus we obtain a new chart Γ_2 in B . Since Γ_2 is included in $B - A$, a Lefschetz fibration $f_2: M_2 \rightarrow B$ corresponding to Γ_2 is a fiber sum of a Lefschetz fibration $f_3: M_3 \rightarrow S^2$ with a trivial Σ_g -bundle over B . Drawing a copy of Γ_2 in S^2 , we have a chart Γ_3 corresponding to f_3 . The



Figure 11: Chart Γ_b 

Figure 12: Channel change




Figure 13: Channel change

signature of a Lefschetz fibration over S^2 described by a free edge is equal to zero because $\tau_g(A, A^{-1}) = 0$ for any $A \in \text{Sp}(2g, \mathbb{Z})$ (see Meyer [30, Section 2]). Hence we have

$$\sigma(M) = \sigma(M_1) = \sigma(M_2) = \sigma(M_3) + \sigma(\Sigma_g \times B) = \sigma(M_3)$$

by Theorem 2.8 and Novikov's additivity. Since we did not change the numbers of white vertices and black vertices of type $\ell_2(h)$ to make Γ_3 from Γ , we see $\sigma(\Gamma_3) = \sigma(\Gamma)$. Therefore we conclude that $\sigma(M) = \sigma(\Gamma)$ because we have already shown that $\sigma(M_3) = \sigma(\Gamma_3)$ in Case 1. $\square$

4 Stabilization theorems

In this section we prove two theorems on stabilization of Lefschetz fibrations under taking fiber sums with copies of a fixed Lefschetz fibration.

Following Auroux [2], we first introduce a notion of universality for Lefschetz fibrations. Suppose that g is greater than two.

Definition 4.1 A Lefschetz fibration of genus g over S^2 is called *universal* if it is irreducible, chiral, and it contains $2g + 1$ singular fibers of type I^+ whose vanishing cycles $a_0, a_1, \dots, a_{2g} \subset \Sigma_g$ satisfies the following conditions: (i) a_i and a_{i+1} intersect transversely at one point for every $i \in \{1, \dots, 2g - 1\}$; (ii) a_0 and a_4 intersect transversely at one point; (iii) a_i and a_j does not intersect for other pairs (i, j) . A Lefschetz fibration over S^2 is universal if and only if it is described by a chart Γ_0 depicted in Figure 14 by virtue of Proposition 2.7, where the blank labeled with T_0 is filled only with edges, white vertices, and black vertices of type $\ell_1(i)$.




Figure 14: Universal chart Γ_0

Remark 4 A universal Lefschetz fibration does exist for every g greater than two. For example, Lefschetz fibrations $f_g^0, f_g^A, f_g^B, f_g^C, f_g^D$ constructed by Auroux [2] are universal except f_g^D for $g = 3$. There would be many universal Lefschetz fibrations of genus g for a fixed g .

We now state the first of our main theorems. Let B be a connected closed oriented surface and $f_0: M_0 \rightarrow S^2$ a universal Lefschetz fibration of genus g .

Theorem 4.2 Let $f: M \rightarrow B$ and $f': M' \rightarrow B$ be Lefschetz fibrations of genus g . There exists a non-negative integer N such that $f \# Nf_0$ is isomorphic to $f' \# Nf_0$ if and only if the following conditions hold: (i) $n_0^\pm(f) = n_0^\pm(f')$; (ii) $n_h^\pm(f) = n_h^\pm(f')$ for every $h = 1, \dots, [g/2]$; (iii) $\sigma(M) = \sigma(M')$.

Remark 5 Auroux [2] proved the ‘if’ part of Theorem 4.2 for chiral Lefschetz fibrations over S^2 under the assumption that f and f' have sections with the same self-intersection number. Hasegawa [13] removed the assumption about existence and self-intersection number of sections in Auroux’s theorem by using chart description.

Remark 6 The isomorphism class of a fiber sum $f \#_{\Psi} f_0$ of a Lefschetz fibration f with a universal Lefschetz fibration f_0 does not depend on a choice of an orientation preserving diffeomorphism Ψ (see Proof of Theorem 4.2).

Proof of Theorem 4.2 We first prove the ‘if’ part. Assume that f and f' satisfy the conditions (i), (ii), and (iii). Let Γ and Γ' be charts in B corresponding to f and f' , respectively. We suppose that f_0 is described by a chart Γ_0 depicted in Figure 14. Since every edge has two adjacent vertices, the sum of the signed numbers of adjacent edges for all vertices of Γ is equal to zero:

$$10n_3(\Gamma) + n_4(\Gamma) - \sum_{i=0}^{2g} m_1(i)(\Gamma) - 4 \sum_{h=1}^{[g/2]} h(2h+1) \cdot m_2(h)(\Gamma) = 0.$$

A similar equality for Γ' also holds. Interpreting the conditions (i) and (ii) as conditions on Γ and Γ' , we have $\sum_{i=0}^{2g} m_1(i)(\Gamma) = \sum_{i=0}^{2g} m_1(i)(\Gamma')$ and $m_2(h)(\Gamma) = m_2(h)(\Gamma')$ for $h = 1, \dots, [g/2]$. Thus we obtain

$$10n_3(\Gamma) + n_4(\Gamma) = 10n_3(\Gamma') + n_4(\Gamma').$$

On the other hand, we have

$$-6n_3(\Gamma) - n_4(\Gamma) = -6n_3(\Gamma') - n_4(\Gamma')$$

by the condition (iii), Theorem 3.5, and $m_2(h)(\Gamma) = m_2(h)(\Gamma')$ for $h = 1, \dots, [g/2]$. Hence $n_3(\Gamma) = n_3(\Gamma')$ and $n_4(\Gamma) = n_4(\Gamma')$.

Let N be an integer larger than both of the number of edges of Γ and that of Γ' . Choose a base point $b_0 \in B - (\Gamma \cup \Gamma')$. The fiber sum $f \# N f_0$ is described by a chart $(\dots((\Gamma \#_{w_1} \Gamma_0) \#_{w_2} \Gamma_0) \dots) \#_{w_N} \Gamma_0$ for some words $w_1, \dots, w_N$ in $\mathcal{X} \cup \mathcal{X}^{-1}$. Since hoops surrounding Γ_0 can be removed by use of the edges of Γ_0 as in Figure 15, the chart is transformed into a product $\Gamma \oplus N\Gamma_0$ by channel changes. Similarly, the fiber sum $f' \# N f_0$ is described by a product $\Gamma' \oplus N\Gamma_0$.




Figure 15: Removing a hoop

We choose and fix $2g + 1$ edges of Γ_0 which are labeled with $0, 1, \dots, 2g$ and adjacent to black vertices. We apply chart moves only to these edges in the following. Since Γ_0 can pass through any edge of Γ as shown in Figure 16, we can move Γ_0 to any region of $B - \Gamma$ by channel changes. For each edge of Γ , we move a copy of Γ_0 to a region




Figure 16: Passing through an edge

adjacent to the edge and apply a channel change to the edge and Γ_0 as in Figure 16 (a) and (b). Applying chart moves of transition to each component of the chart as in Figure 17, we remove white vertices of type $r_1(i, j)^{\pm 1}, r_2(i)^{\pm 1}, r_5^{\pm 1}$ to obtain a union of copies of $L_1(i), L_2(h), L_2(h)^*, R_3, R_3^*, R_4, R_4^*, \Gamma_0$ shown in Figure 18, 19, 20, where we use a simplification of diagrams as in Figure 21.




Figure 17: Chart moves of transition

If there is a pair of R_3 and R_3^* , we remove them by a death of a pair of white vertices to obtain many copies of Γ_0 . Similarly, we remove a pair of R_4 and R_4^* . Since there is at least one Γ_0 , any copy of $L_1(i)$ can be transformed into $L_1(1)$ as in Figure 22.





Figure 18: Charts $L_1(i)$, $L_2(h)$, $L_2(h)^*$ Figure 19: Charts R_3 and R_3^* Figure 20: Charts R_4 and R_4^* 

Figure 21: Simplification of diagram

Thus we have a union Γ_1 of $m_1^-(\Gamma)$ copies of $L_1(1)$, $m_2^+(h)(\Gamma)$ copies of $L_2(h)$, $m_2^-(h)(\Gamma)$ copies of $L_2(h)^{-1}$, $|n_3(\Gamma)|$ copies of R_3 (or R_3^*), $|n_4(\Gamma)|$ copies of R_4 (or R_4^*), and k copies of Γ_0 for some k . A similar argument implies that $\Gamma' \oplus N\Gamma_0$ is transformed into a union Γ'_1 of $m_1^-(\Gamma')$ copies of $L_1(1)$, $m_2^+(h)(\Gamma')$ copies of $L_2(h)$, $m_2^-(h)(\Gamma')$ copies of $L_2(h)^{-1}$, $|n_3(\Gamma')|$ copies of R_3 (or R_3^*), $|n_4(\Gamma')|$ copies of R_4 (or R_4^*), and k' copies of Γ_0 for some k' by chart moves of type W and chart moves of transition. By virtue of the conditions (i) and (ii) together with $n_3(\Gamma) = n_3(\Gamma')$, $n_4(\Gamma) = n_4(\Gamma')$, $m_1^+(\Gamma \oplus N\Gamma_0) = m_1^+(\Gamma_1)$, and $m_1^+(\Gamma' \oplus N\Gamma_0) = m_1^+(\Gamma'_1)$, we conclude that $k = k'$ because of $m_1^+(\Gamma_0) \neq 0$. Hence Γ_1 is transformed into Γ'_1 by an ambient isotopy of B relative to b_0 , which means that $\Gamma \oplus N\Gamma_0$ is transformed into $\Gamma' \oplus N\Gamma_0$ by


Figure 22: Changing a label ($j = i + 1$ or $(i, j) = (4, 0)$)

chart moves of type W, chart moves of transition, and ambient isotopies of B relative to b_0 . Therefore $f\#Nf_0$ is (strictly) isomorphic to $f'\#Nf_0$ by Theorem 2.7.

We next prove the ‘only if’ part. Take a non-negative integer N so that $f\#Nf_0$ is isomorphic to $f'\#Nf_0$. Since an isomorphism preserves numbers and types of vanishing cycles and signatures, we have $n_0^\pm(f\#Nf_0) = n_0^\pm(f'\#Nf_0)$, $n_h^\pm(f\#Nf_0) = n_h^\pm(f'\#Nf_0)$ for every $h = 1, \dots, [g/2]$, and $\sigma(M\#_F NM_0) = \sigma(M'\#_F NM_0)$. The conditions (i), (ii), (iii) follows from additivity of $n_0^\pm, n_h^\pm, \sigma$ under fiber sum. $\square$

Definition 4.3 A Lefschetz fibration of genus g over S^2 is called *elementary* if it contains exactly two singular fibers of type I^+ and of type I^- which have the same vanishing cycles. A chart $L_1(i)$ in S^2 corresponds to an elementary Lefschetz fibration.

Remark 7 Two elementary Lefschetz fibrations of genus g are isomorphic to each other. The total space of an elementary Lefschetz fibration of genus g is diffeomorphic to $\Sigma_{g-1} \times S^2 \# S^1 \times S^3$.

We state the second of our main theorems. Let B be a connected closed oriented surface and $f_\star: M_\star \rightarrow S^2$ an elementary Lefschetz fibration of genus g .

Theorem 4.4 Let $f: M \rightarrow B$ and $f': M' \rightarrow B$ be Lefschetz fibrations of genus g . There exists a non-negative integer N such that a fiber sum $f\#Nf_\star$ is isomorphic to a fiber sum $f'\#Nf_\star$ if and only if the following conditions hold: (i) $n_0^\pm(f) = n_0^\pm(f')$; (ii) $n_h^\pm(f) = n_h^\pm(f')$ for every $h = 1, \dots, [g/2]$; (iii) $\sigma(M) = \sigma(M')$.

Remark 8 In contrast to Theorem 4.2, the isomorphism class of a fiber sum $f \#_{\Psi} f_{\star}$ of a Lefschetz fibration f with an elementary Lefschetz fibration $f_{\star}$ depends on a choice of an orientation preserving diffeomorphism Ψ in general.

Proof of Theorem 4.4 We only show the ‘if’ part. The ‘only if’ part is the same as that of the proof of Theorem 4.2.

Assume that f and f' satisfy the conditions (i), (ii), and (iii). Let Γ and Γ' be charts in B corresponding to f and f' , respectively. It follows from the same argument as in the proof of Theorem 4.2 that $n_3(\Gamma) = n_3(\Gamma')$ and $n_4(\Gamma) = n_4(\Gamma')$. Let N be an integer larger than both of the number of edges of Γ and that of Γ' . Choose a base point $b_0 \in B - (\Gamma \cup \Gamma')$ and a disk D in $B - \Gamma$ centered at b_0 . We denote the set of edges of Γ by $E(\Gamma)$. For each $e \in E(\Gamma)$, we choose a point b_e in a region of $B - \Gamma$ adjacent to e , and a simple path γ_e from b_e to b_0 which intersects with edges of Γ transversely and does not intersect with vertices of Γ . Let w_e be the intersection word of γ_e with respect to Γ and $i_e \in \{0, 1, \dots, 2g\}$ the label of e . We choose a family $\{D_e\}_{e \in E(\Gamma)}$ of disjoint disks included in D and put the chart Γ_e depicted in Figure 23 in D_e for each e . Taking the union of Γ with Γ_e for all $e \in E(\Gamma)$ and with $N - \#E(\Gamma)$ copies of $L_1(1)$, we obtain a new chart Γ_1 in B , which describes a fiber sum $f \# N f_{\star}$.




Figure 23: Chart Γ_e

For each edge e of Γ , we apply channel changes as in Figure 24 to let a free edge pass through the edges intersecting with γ_e . We then apply a channel change as in Figure 25 to ‘cut’ e into two edges.

Applying deaths of pairs of white vertices appropriately, we obtain a union Γ_2 of $m_2^+(h)(\Gamma)$ copies of $\tilde{L}_2(h)$, $m_2^-(h)(\Gamma)$ copies of $\tilde{L}_2(h)^*$, $|n_3(\Gamma)|$ copies of $\tilde{R}_3$ (or $\tilde{R}_3^*$), $|n_4(\Gamma)|$ copies of $\tilde{R}_4$ (or $\tilde{R}_4^*$), and k_i copies of $L_1(i)$ for some k_i , where $\tilde{R}_3$ and $\tilde{R}_4$ are charts depicted in Figure 26, $\tilde{L}_2(h)^* = L_2(h)^*$, and $\tilde{L}_2(h)$ (resp. $\tilde{R}_3^*$, $\tilde{R}_4^*$) is the mirror image of $\tilde{L}_2(h)^*$ (resp. $\tilde{R}_3$, $\tilde{R}_4$) with edges orientation reversed. Similarly, Γ' is transformed into Γ'_1 , which describes a fiber sum $f' \# N f_{\star}$, and then a union Γ'_2 of $m_2^+(h)(\Gamma')$ copies of $\tilde{L}_2(h)$, $m_2^-(h)(\Gamma')$ copies of $\tilde{L}_2(h)^*$, $|n_3(\Gamma')|$ copies of $\tilde{R}_3$ (or $\tilde{R}_3^*$), $|n_4(\Gamma')|$ copies of $\tilde{R}_4$ (or $\tilde{R}_4^*$), and k'_i copies of $L_1(i)$ for some k'_i .




Figure 24: Channel change





Figure 25: Channel change

Figure 26: Charts $\tilde{R}_3$ and $\tilde{R}_4$

A similar argument on the number m_1^+ as in the proof of Theorem 4.2 implies that $k_0 + k_1 + \cdots + k_{2g} = k'_0 + k'_1 + \cdots + k'_{2g}$. Adding $|k_i - k'_i|$ copies of $L_1(i)$ to either Γ_2 or Γ'_2 if necessary, we may assume that $k_i = k'_i$ for every $i \in \{0, 1, \dots, 2g\}$. Hence Γ_2 is transformed into Γ'_2 by an ambient isotopy of B relative to b_0 , which means that $f\#Nf_*$ is (strictly) isomorphic to $f'\#Nf_*$ by Theorem 2.7. $\square$

Let g be an integer greater than two and $B_1, \dots, B_r$ connected closed oriented surfaces. We consider a Lefschetz fibration $f_i: M_i \rightarrow B_i$ of genus g for each $i \in \{1, \dots, r\}$, and a universal Lefschetz fibration $f_0: M_0 \rightarrow S^2$ of genus g .

Proposition 4.5 *For (possibly different) fiber sums f and f' of $f_1, \dots, f_r$, fiber sums $f\#f_0$ and $f'\#f_0$ are isomorphic to each other.*

Proof Let Γ and Γ' be charts corresponding to f and f' . Since hoops surrounding a component of Γ (and Γ') can be removed by use of the edges of Γ_0 as in Figure 15, $\Gamma\#\Gamma_0$ and $\Gamma'\#\Gamma_0$ are transformed into the same chart. $\square$

Remark 9 Proposition 4.5 implies that there are many examples of non-isomorphic Lefschetz fibrations with the same base, the same fiber, and the same numbers of singular fibers of each type which become isomorphic after one stabilization. For example, the Lefschetz fibration on $E(n)_K$ constructed by Fintushel and Stern [8, Theorem 14] (see also Park and Yun [33]) for a fibered knot K becomes isomorphic to that on $E(n)_{K'}$ for another fibered knot K' of the same genus after one stabilization. Similar results hold for Lefschetz fibrations on $Y(n; K_1, K_2)$ constructed by Fintushel and Stern [8, § 7] (see also Park and Yun [34]) as well as fiber sums of (generalizations of) Matsumoto's fibration studied by Ozbagci and Stipsicz [32], Korkmaz [22, 23], and Okamori [31].

5 Variations and problems

In this section we discuss possible variations of chart description for Lefschetz fibrations.

If we replace the triple $(\mathcal{X}, \mathcal{R}, \mathcal{S})$ defined in Section 2 with other triples, we obtain various chart descriptions for Lefschetz fibrations (see Kamada [18]).

We first choose large $\mathcal{X}$, $\mathcal{R}$, and $\mathcal{S}$. Let $\mathcal{X}$ be the set of right-handed Dehn twists along simple closed curves in Σ_g and $\mathcal{S}$ the set of Dehn twists along non-trivial simple closed curves in Σ_g . By virtue of a theorem of Luo [26], $\langle \mathcal{X} | \mathcal{R} \rangle$ gives an infinite presentation of $\mathcal{M}_g$ for the set $\mathcal{R}$ of the following four kind of words: (0) trivial relator $A := a$, where a is the Dehn twist along trivial simple closed curve on Σ_g ; (1) braid relator $T := b^{-1}abc^{-1}$, where $a, b, c \in \mathcal{X} \cup \mathcal{X}^{-1}$ and the curve for c is the image of the curve for a by b ; (2) chain relator $C := (c_2c_1)^6d^{-1}$, where $c_1, c_2, d \in \mathcal{X} \cup \mathcal{X}^{-1}$ and the curves for c_1 and c_2 intersect transversely at one point and the curve for d is the boundary curve of a regular neighborhood of the union of the curves for c_1 and c_2 ; (3) lantern relator $L := cbad_4^{-1}d_3^{-1}d_2^{-1}d_1^{-1}$, where the curves for a and b intersect transversely at two points with algebraic intersection number zero, the curve

for c is obtained by resolving the intersections of these two curves, and the curves for d_1, d_2, d_3, d_4 are the boundary curves of a regular neighborhood of those for a, b, c .

Let B be a connected closed oriented surface. Charts in B for the triple $(\mathcal{X}, \mathcal{R}, \mathcal{S})$ defined above have white vertices of type $A^{\pm 1}, T^{\pm 1}, C^{\pm 1}, L^{\pm 1}$ (see Figure 27). For a chart Γ in B , we denote the number of white vertices of type r minus the number of white vertices of type r^{-1} included in Γ by $n_r(\Gamma)$.

Proposition 5.1 *The signature $\sigma(M)$ of the total space M of a Lefschetz fibration $f: M \rightarrow B$ described by Γ is equal to $-n_A(\Gamma) - 7n_C(\Gamma) + n_L(\Gamma)$.*

Proof A similar argument to the proof of Theorem 3.5. □




Figure 27: Vertices of type A, T, C, L

Problem 5.2 Study various properties of Lefschetz fibrations by using chart description for the triple $(\mathcal{X}, \mathcal{R}, \mathcal{S})$ defined above.

We next mention chart description for Lefschetz fibrations with bordered base and fiber. Kamada [18] gave a general theory for charts in a compact oriented surface with boundary. Various presentations of mapping class groups of surfaces with boundary have been investigated by researchers including Gervais [10], Labruère and Paris [25], Margalit and McCammond [27]. Combining these two kinds of studies, one can immediately obtain a chart description for Lefschetz fibrations with bordered base and fiber.

Problem 5.3 Make use of chart description to study PALFs and Stein surfaces.

It would be worth considering compositions of monodromy representations with appropriate homomorphisms and charts corresponding to the compositions. For example, Hasegawa [12, 13] adopted a homomorphism from the m -string braid group B_m to the semi-direct product $(\mathbb{Z}_2)^m \times S_m$, while Endo and Kamada [6] used a standard epimorphism from the hyperelliptic mapping class group of a closed oriented surface of genus g to the mapping class group of a sphere with $2g + 2$ marked points.

References

- [1] D. Auroux, *Fiber sums of genus 2 Lefschetz fibrations*, Turkish J. Math. **27** (2003) 1–10.
- [2] D. Auroux, *A stable classification of Lefschetz fibrations*, Geom. Topol. **9** (2005), 203–217.
- [3] J. Barge and É. Ghys, *Cocycles d’Euler et de Maslov*, Math. Ann. **294** (1992), 235–265.
- [4] R. I. Baykur and S. Kamada, *Classification of broken Lefschetz fibrations with small fiber genera*, arXiv:1010.5814.
- [5] H. Endo and S. Kamada, *Chart description for hyperelliptic Lefschetz fibrations and their stabilization*, arXiv:1306.2707, to appear in Topology Appl.
- [6] H. Endo and S. Kamada, *Counting Dirac braids and hyperelliptic Lefschetz fibrations*, in preparation.
- [7] H. Endo and S. Nagami, *Signature of relations in mapping class groups and non-holomorphic Lefschetz fibrations*, Trans. Amer. Math. Soc. **357** (2004), 3179–3199.
- [8] R. Fintushel and R. J. Stern, *Families of simply connected 4-manifolds with the same Seiberg-Witten invariants*, Topology **43** (2004), 1449–1467.
- [9] J.-M. Gambaudo and É. Ghys, *Braids and signatures*, Bull. Soc. Math. France **133** (2005), 541–579.
- [10] S. Gervais, *A presentation of the mapping class group of a punctured surface*, Topology **40** (2001), 703–725.
- [11] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*, Grad. Stud. Math. **20**, Amer. Math. Soc., Providence, RI, 1999.
- [12] I. Hasegawa, *A certain linear representation of the classical braid group and its application to surface braids*, Math. Proc. Camb. Phil. Soc. **141** (2006), 287–301.
- [13] I. Hasegawa, *Chart descriptions of monodromy representations on oriented closed surfaces*, thesis, Univ. of Tokyo, March 23, 2006.
- [14] K. Hayano, *On genus–1 simplified broken Lefschetz fibrations*, Algebr. Geom. Topol. **11** (2011), 1267–1322.
- [15] S. Kamada, *Surfaces in R^4 of braid index three are ribbon*, J. Knot Theory Ramifications **1** (1992), 137–160.
- [16] S. Kamada, *An observation of surface braids via chart description*, J. Knot Theory Ramifications **4** (1996), 517–529.
- [17] S. Kamada, *Braid and knot theory in dimension four*, Math. Surveys Monogr. **95**, Amer. Math. Soc., Providence, RI, 2002.
- [18] S. Kamada, *Graphic descriptions of monodromy representations*, Topology Appl. **154** (2007), 1430–1446.

- [19] S. Kamada, *Chart description for genus-two Lefschetz fibrations and a theorem on their stabilization*, Topology Appl. **159** (2012), 1041–1051.
- [20] S. Kamada, Y. Matsumoto, T. Matumoto and K. Waki, *Chart description and a new proof of the classification theorem of genus one Lefschetz fibrations*, J. Math. Soc. Japan, **57** (2005), 537–555.
- [21] A. Kas, *On the handlebody decomposition associated to a Lefschetz fibration*, Pacific J. Math. **89** (1980), 89–104.
- [22] M. Korkmaz, *Noncomplex smooth 4-manifolds with Lefschetz fibrations*, Int. Math. Res. Not. IMRN, Vol. 2001, pp. 115–128. .
- [23] M. Korkmaz, *Lefschetz fibrations and an invariant of finitely presentable groups*, Int. Math. Res. Not. IMRN, Vol. 2009, pp. 1547–1572.
- [24] Y. Kuno, *Meyer functions and the signature of fibered 4-manifolds*, arXiv:1204.1701.
- [25] C. Labruère and L. Paris, *Presentations for the punctured mapping class groups in terms of Artin groups*, Algebr. Geom. Topol. **1** (2001), 73–114.
- [26] F. Luo, *A presentation of the mapping class groups*, Math. Res. Lett. **4** (1997), 725–739.
- [27] D. Margalit and J. McCammond, *Geometric presentations for the pure braid group*, J. Knot Theory Ramifications **18** (2009), 1–20.
- [28] Y. Matsumoto, *Diffeomorphism types of elliptic surfaces*, Topology **25** (1986), 549–563.
- [29] Y. Matsumoto, *Lefschetz fibrations of genus two — A topological approach*, in “Topology and Teichmüller spaces” S. Kojima et al, eds., Proc. the 37-th Taniguchi Sympo., pp. 123–148, World Scientific Publishing, River Edge, NJ, 1996.
- [30] W. Meyer, *Die Signatur von Flächenbündeln*, Math. Ann. **201** (1973), 239–264.
- [31] K. Okamori, *A genus 2 Lefschetz fibration on an exotic $\mathbb{CP}^2 \# 9\overline{\mathbb{CP}^2}$* , master’s thesis (in Japanese), Osaka University, February 2011.
- [32] B. Ozbagci and A. I. Stipsicz, *Noncomplex smooth 4-manifolds with genus-2 Lefschetz fibrations*, Proc. Amer. Math. Soc. **128** (2000), 3125–3128.
- [33] J. Park and K.-H. Yun, *Nonisomorphic Lefschetz fibrations on knot surgery 4-manifolds*, Math. Ann. **345** (2009), 581–597.
- [34] J. Park and K.-H. Yun, *Lefschetz fibration structure on knot surgery 4-manifolds*, Michigan Math. J. **60** (2011), 525–544.
- [35] V. G. Turaev, *First symplectic Chern class and Maslov indices*, J. Soviet Math. **37** (1987), 1115–1127.
- [36] B. Wajnryb, *A simple presentation for the mapping class group of an orientable surface*, Israel J. Math. **45** (1983), 157–174.
- [37] B. Wajnryb, *An elementary approach to the mapping class group of a surface*, Geom. Topol. **3** (1999), 405–466.

- [38] C. T. C. Wall, *Non-additivity of the signature*, Invent. Math. 7 (1969), 269–274.

Department of Mathematics, Tokyo Institute of Technology
2-12-1 Oh-okayama, Meguro-ku, Tokyo 152-8551, Japan

Ministry of Health, Labour and Welfare
1-2-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-8916, Japan

Department of Mathematics, Osaka City University
3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

endo@math.titech.ac.jp, hasegawa-isao01@mhlw.go.jp,
skamada@sci.osaka-cu.ac.jp